

## A generalization of product inequality for the higher topological complexity

Nguyen Viet Dung (Hanoi, Vietnam)

Nguyen Van Ninh (Thai Nguyen, Vietnam)

(Received Oct. 22, 2022)

**Abstract.** In [4] M. Farber defined the topological complexity  $TC(X)$  of a path-connected space  $X$ . Generalizing this notion, ten years later, Yu. Rudyak introduced a sequence of invariants, called the higher topological complexities  $TC_n(X)$ , for any path-connected space  $X$  in [7]. These invariants have their origin in the notion of the Schwarz genus of a fibration defined in [8]. One of the tools used to calculate these invariants is the product inequality for the Schwarz genus. In this paper, we will give a generalization of the product inequality of the higher topological complexity.

### 1. Introduction

Let  $X$  be a path-connected topological space,  $PX$  the space of all continuous paths  $\gamma : I = [0, 1] \rightarrow X$  with the usual compact-open topology.

Let's consider the map

$$\begin{aligned} \pi : PX &\longrightarrow X \times X \\ \gamma &\longmapsto (\gamma(0), \gamma(1)) \end{aligned}$$

---

*Key words and phrases:* higher topological complexity, configuration space, Schwarz genus, product inequality.

*2020 Mathematics Subject Classification:* 55R80, 55M30

The first author is supported by the grant NCVCC01.16/22-22

In [4], M. Farber defined the topological complexity  $TC(X)$  of  $X$  as the smallest number  $k$  such that there exists an open covering  $\{U_i, i = 1, \dots, k\}$  of  $X \times X$  with a continuous section  $s_i : U_i \rightarrow PX$  of  $\pi$  on each  $U_i$ , i.e.  $\pi \circ s_i = id_{U_i}$ .

In 2010 Yu. Rudyak (see [7]) introduced a series of invariants, denoted by  $TC_n(X)$ ,  $n \geq 2$ , for any path-connected space  $X$ .  $TC_n(X)$  is called the higher topological complexity of  $X$ . It coincides with the topological complexity  $TC(X)$  defined by M. Farber when  $n = 2$ .

Since then, the higher topological complexity has been computed for many topological spaces like spheres in [7] and product of spheres in [1], wedge sum of spheres of different dimensions in [3], configuration spaces on Euclidean spaces in [5], configuration space of distinct ordered points on compact Riemann surfaces of genus  $g$  in [6], the complement of some classes of complex hyperplane arrangements in [2].

Being closely related to the notion of the Schwarz genus and the Lusternik-Shnirelman category, the higher topological complexity inherited many interesting properties of these invariants. Some of these properties such as some evaluations from above or from below are used in computing the higher topological complexity  $TC_n$ .

One of the important evaluations of higher topological complexity is the product inequality saying that if  $X$  and  $Y$  are path-connected spaces, then

$$TC_n(X \times Y) \leq TC_n(X) + TC_n(Y) - 1.$$

In this paper, we give a generalization of this inequality. The paper is organized as follows. In section 2 we investigate the higher topological complexity and some of its properties. We formulate and prove our main result in section 3.

## 2. Higher topological complexity

For  $n = 2, 3, \dots$  let  $J_n$  denote the wedge sum of  $n$  closed unit intervals  $[0, 1]_i$ ,  $i = 1, \dots, n$  with 0 as attached point. Suppose that  $X^{J_n}$  denotes the space of all continuous maps  $\gamma : J_n \rightarrow X$  with compact-open topology. Consider the map

$$\begin{aligned} e_n^X : X^{J_n} &\longrightarrow X^n, \\ \gamma &\longmapsto (\gamma(1_1), \dots, \gamma(1_n)) \end{aligned}$$

where  $1_i$  is the unit in  $[0, 1]_i$  respectively.

**Definition 2.1** (see [7]). *The higher topological complexity  $TC_n(X)$  of the space  $X$  is the smallest number  $k$  such that there is an open covering  $\{U_i, i = 1, \dots, k\}$  of  $X^n$  and there exists a continuous section  $s_i : U_i \rightarrow X^{J_n}$  of  $e_n^X$ , on each  $U_i$ , i.e.,  $e_n^X \circ s_i = id_{U_i}$ .*

Obviously, when  $n = 2$ ,  $TC_2(X)$  coincides the topological complexity  $TC(X)$  defined by M.Faber.

**Remark 2.1.** *It is known that the map  $e_n^X$  is a fibration in the sense of Serre. By definition, the higher topological complexity  $TC_n(X)$  of the space  $X$  is exactly the Schwarz genus of the fibration  $e_n^X$  (see [8]). Moreover, as it is indicated in [7],  $e_n^X$  is a fibrational substitute of the diagonal map  $d_n$ , i.e. there exists a homotopy equivalence  $h : X \rightarrow X^{J_n}$  such that  $d_n = e_n^X \circ h$ . Therefore,  $TC_n(X)$  is also called the Schwarz genus of  $d_n$ .*

Similar to the topological complexity, the higher topological complexity  $TC_n(X)$  is a homotopy invariant. This property has been proved for the topological complexity  $TC(X)$  in [4]. We present here proof of this important property for the  $TC_n(X)$ .

**Proposition 2.1.** *Suppose that  $X$  is homotopic to  $Y$ . Then  $TC_n(X) = TC_n(Y)$ .*

**Proof.** Assume that there exist continuous maps  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  such that  $f \circ g \simeq id_Y$ . We will prove that  $TC_n(Y) \leq TC_n(X)$ .

Let  $U$  be a open set in  $X^n$  such that there exists a continuous section  $s : U \rightarrow X^{J_n}$  of  $e_n^X$ .

For  $(B_1, \dots, B_n) \in U$ , we have  $s(B_1, \dots, B_n)$  is a map  $\gamma : J_n \rightarrow X$ . For any  $i = 1, \dots, n$ , let  $\gamma_i$  be the path in  $X$  defined by  $\gamma_i(t) = \gamma|_{[0,1]_i}(t)$ , where  $[0, 1]_i$  denotes the  $i^{th}$  unit interval in the wedge  $J_n = [0, 1] \vee \dots \vee [0, 1]$ .

Let's consider the set

$$V = (g \times \dots \times g)^{-1}(U) = \{(A_1, \dots, A_n) \in Y^n | (g(A_1), \dots, g(A_n)) \in U\}.$$

We are going to construct a continuous section  $\sigma : V \rightarrow Y^{J_n}$  of  $e_n^Y$  on this open set  $V$  of  $Y^n$ .

Suppose that  $H_t : Y \rightarrow Y$  is the homotopy  $id_Y \simeq f \circ g$  with  $H_0 = id_Y$ ,  $H_1 = f \circ g$ . For  $(A_1, A_2, \dots, A_n) \in V$  we have  $(g(A_1), \dots, g(A_n)) \in U$  and therefore there exists a continuous section  $s$  of  $e_n^X$ . As mentioned above,  $s(g(A_1), \dots, g(A_n))$  is a path  $\gamma : J_n \rightarrow X^n$ . Now, we define  $[A_1, A_i]$  to be the path in  $Y$  connecting  $A_1$  to  $A_i$ ,  $i = 1, \dots, n$ , by

$$[A_1, A_i](t) = \begin{cases} H_{3t}(A_1), & \text{if } 0 \leq t < \frac{1}{3} \\ f(s_{(g(A_1), g(A_i))}(3t - 1)), & \text{if } \frac{1}{3} \leq t < \frac{2}{3} \\ H_{3(1-t)}(A_i), & \text{if } \frac{2}{3} \leq t \leq 1 \end{cases}$$

Here  $s_{(g(A_1), g(A_i))}$  denotes the path in  $X$ , connecting  $g(A_1)$  to  $g(A_i)$ , defined from  $s$  by

$$s_{(g(A_1), g(A_i))}(t) = \begin{cases} \gamma_1(1 - 2t), & \text{if } 0 \leq t < \frac{1}{2} \\ \gamma_i(2t - 1), & \text{if } \frac{1}{2} \leq t \leq 1 \end{cases}.$$

The section  $\sigma$  of  $e_n^Y$  on  $V$  is defined by putting

$$\sigma(A_1, \dots, A_n) = ([A_1, A_1], [A_1, A_2], \dots, [A_1, A_n]).$$

Here the right hand side denotes a map from  $J^n$  to  $X^n$  having its restriction on the  $i^{\text{th}}$  summand  $[0, 1]_i$  of  $J^n$  to be the path  $[A_1, A_i]$ . It is easy to see that  $\sigma$  defined above is continuous and  $e_n^Y \circ \sigma = id_V$ . Thus,  $TC_n(Y) \leq TC_n(X)$ .

Similar arguments will prove  $TC_n(X) \leq TC_n(Y)$ . And these imply the Proposition.

The following property shows the relation between the higher topological complexity  $TC_n(X)$  of a space  $X$  and homotopy properties of  $X$ .

**Proposition 2.2.** *If  $X$  is a finite  $r$ -connected polyhedron, then*

$$TC_n(X) < \frac{n \dim X + 1}{r + 1} + 1.$$

*In particular  $r = 0$  (i.e.  $X$  is path-connected) then*

$$TC_n(X) \leq n \dim X + 1.$$

**Proof.** As it is mentioned in the Remark after Definition 2.1, the higher topological complexity  $TC_n(X)$  is nothing but the Schwarz genus of the fibration  $e_n^X$ . The proposition is a consequence of the similar property of the Schwarz genus (see [8, Theorem 5]).

A lower bound for the higher topological complexity of the space  $X$  is given in terms of its cohomology with coefficient in any field  $\mathbb{K}$ .

**Definition 2.2.** *Let  $X$  be a finite path-connected polyhedron. Suppose that  $n$  is an integer,  $d_n$  is the diagonal map  $d_n : X \rightarrow X^n$  and  $\mathbb{K}$  is a field.*

1. *The kernel of the homomorphism  $d_n^* : H^*(X^n; \mathbb{K}) \rightarrow H^*(X; \mathbb{K})$  is called the  $n$ -zero divisor of  $X$ .*
2.  *$d_n$ -zero divisor cup length of  $X$ , denoted by  $cl(X, n)$  (see[1]), is the maximal number  $k$  such that there exists  $k$  elements of the  $n$ -zero divisor of  $X$  satisfying  $u_1 \cup u_2 \dots \cup u_k \neq 0$ .*

Then  $cl(X, n)$  will be a lower bound of  $TC_n(X)$ . Precisely, we have the following proposition, which follows from [8, Theorem 4]. Detailed proofs can be found in [7] and [1].

**Proposition 2.3.** *Suppose that  $n$  is an integer,  $n \geq 2$ .*

*i) Let  $X$  be a path-connected topology space. Then*

$$TC_n(X) \geq cl(X, n) + 1.$$

*ii) For finite path-connected polyhedra  $X$  and  $Y$  we have*

$$cl(X \times Y, n) \geq cl(X, n) + cl(Y, n).$$

The next property, usually called product inequality, gives us an estimate of the higher topological complexity of the product space by those of its factors. This inequality is very useful in computing the higher topological complexity of many spaces.

**Proposition 2.4.** *For path-connected spaces  $X$  and  $Y$ . If  $(X \times Y)^n$  is normal, then we have*

$$(2.1) \quad TC_n(X \times Y) \leq TC_n(X) + TC_n(Y) - 1.$$

Similar property for the Schwarz genus is known in [8, proposition 22] and for the topological complexity  $TC(X)$  in [4]. A detailed proof for the case of the higher topological complexity can be found in [1].

### 3. A generalization of inequality product

In this section, we will generalize the inequality product for the higher topological complexity.

**Theorem 3.1.** *Suppose that  $E, X$  are finite path-connected CW-complexes,  $(Y, y_0)$  is a pointed space and  $p : E \rightarrow X$  is a continuous map such that the following conditions hold*

*i) For all  $x \in X$ , the fiber  $p^{-1}(x)$  is homotopic to  $Y$ .*

*ii) The map  $p$  accepts a section  $s : X \rightarrow E$ , i.e.,  $p \circ s = id_X$ .*

*iii) There exists a family of homotopy equivalences  $h_x : p^{-1}(x) \rightarrow Y$  depending continuously on  $x \in X$  such that  $h_x(s(x)) = y_0$ .*

Then,

$$TC_n(E) \leq TC_n(X) + TC_n(Y) - 1.$$

To prove our main result, we first need some technical lemmas.

**Lemma 3.1.** *Let  $Y$  be a finite path-connected CW-complex,  $U$  an open set in  $Y^n$  on which there is a continuous section  $s_U : U \rightarrow Y^{J^n}$  of  $e_n^Y$ . Then, for any  $y_0 \in Y$  there exists always a continuous section  $s'_U : U \rightarrow Y^{J^n}$  of  $e_n^Y$  such that  $s'_U(A_1, \dots, A_n)(0) = y_0$  for any  $(A_1, \dots, A_n) \in U$ .*

**Proof.** We first consider the case when  $y_0 \in U$ .

For  $(A_1, \dots, A_n) \in U$  we have  $s_U(A_1, \dots, A_n)$  is a map  $\gamma : J^n \rightarrow Y^n$ . Denote  $\gamma(0) = P \in Y$ . Define  $\gamma_i$  to be the restriction of  $\gamma$  on the  $i^{\text{th}}$  unit interval of  $J^n$ . That is,  $\gamma_i$  is a path connecting  $P$  and  $A_i$ .

By means of the section  $s_U$ , it implies that for any point  $P \in U$ , there exists a path connecting  $y_0$  to  $P$  and this path depends continuously on  $P$ . Let denote this path by  $\ell_P$ .

Now the section  $s'_U$  is defined by putting  $s'_U(A_1, \dots, A_n)$  to be a map  $\gamma' : J^n \rightarrow Y^n$ , where  $\gamma'$  has its restriction on the  $i^{\text{th}}$  unit interval of  $J^n$  as  $\ell_P * \gamma_i$ . It is easy to see that  $s'_U(A_1, \dots, A_n)$  is a continuous section of  $e_n^Y$  and  $s'_U(A_1, \dots, A_n)(0) = y_0$ .

Now suppose that  $y_0 \notin U$ . Let fix a point  $y_1 \in U$ . By the above arguments, we can construct a continuous section  $\tilde{s}_U$  of  $e_n^Y$  such that  $\tilde{s}_U(A_1, \dots, A_n)(0) = y_1$ . Suppose that  $\tilde{s}_U(A_1, \dots, A_n)$  is a map  $\tilde{\gamma} : J^n \rightarrow Y^n$ . Then its restriction  $\tilde{\gamma}_i$  on the  $i^{\text{th}}$  unit interval of  $J^n$  is a path connecting  $y_1$  to  $A_i$ . Fix a path  $\ell$  in  $Y$  connecting  $y_0$  and  $y_1$ . Now we define the section  $s'$  by putting  $s'_U(A_1, \dots, A_n)$  to be a map  $J^n \rightarrow Y^n$  having its restriction on the  $i^{\text{th}}$  unit interval of  $J^n$  to be  $\ell * \tilde{\gamma}_i$ . Obviously, the defined map  $s'$  is a section of  $e_n^Y$  having  $s'_U(A_1, \dots, A_n)(0) = y_0$ .

The following lemma is implied from Proposition 20 of [8] and the fact that  $TC_n(X)$  coincides with the Schwarz genus of the fibration  $e_n^X$ .

**Lemma 3.2.** *Given the topological space  $X$ , let  $C = \{C_1, \dots, C_p\}$  and  $D = \{D_1, \dots, D_q\}$  be open covering of  $X^n$  such that on each  $C_i \cap D_j$  there exists local section of  $e_n$ . Then*

$$TC_n(X) \leq p + q - 1.$$

**Proof.** [Proof of Theorem 3.1] Suppose that  $TC_n(X) = p$ . By definition, there is an open covering  $U = \{U_1, \dots, U_p\}$  of  $X^n$  such that there exists a section of  $e_n^X$  on each  $U_i$ . Put  $C_i = \{(A_1, \dots, A_n) \in E^n \mid (p(A_1), \dots, p(A_n)) \in U_i\}$ . Then  $C = \{C_1, \dots, C_p\}$  is an open covering of  $E^n$

Suppose that  $TC_n(Y) = q$ , and  $V = \{V_1, \dots, V_q\}$  is an open covering of  $Y^n$  such that there exists a section of  $e_n^Y$  on each  $V_j$ . Let

$$D_j = \{(A_1, \dots, A_n) \in E^n \mid (h_{p(A_1)}(A_1), \dots, h_{p(A_n)}(A_n)) \in V_j\}.$$

Then  $D = \{D_1, \dots, D_q\}$  is an open covering of  $E^n$ . Let us fix a section of  $e_n^Y$  on each  $V_j$  as that in the Lemma 3.1.

We will now construct a section of  $e_n^E$  on each  $C_i \cap D_j$ ,  $i = 1, \dots, p$ ,  $j = 1, \dots, q$ . Suppose that  $(A_1, \dots, A_n) \in C_i \cap D_j \subset E^n$ .

By definition  $(A_1, \dots, A_n) \in C_i$  means that  $(p(A_1), \dots, p(A_n)) \in U_i$ . Since there exists a continuous section  $s_1$  of  $e_n^X$  on  $U_i$ , there is a path going from  $p(A_1)$  to  $p(A_i)$  defined by this section  $s_1$ . Let denote  $\gamma_2$  the image of this path by the section  $s$  of  $p$ .

Since  $(A_1, \dots, A_n) \in D_j$  it implies that  $(h_{p(A_1)}(A_1), \dots, h_{p(A_n)}(A_n)) \in V_j$ . By assumption, there exists a section  $s_2$  of  $e_n^Y$  on this  $V_j$ . Let's choose  $s_2$  to be the one defined in the Lemma 3.1. This section  $s_2$  defines a path in  $Y$  connecting  $h_{p(A_1)}(A_1)$  to the point  $y_0$ . We denote by  $\gamma_1$  the inverse image of this path by the homotopy equivalence  $h_{p(A_1)}$ .

Similarly, the section  $s_2$  defines a path in  $Y$  connecting  $y_0$  to the point  $h_{p(A_i)}(A_i)$ ,  $i = 1, \dots, n$ . We denote by  $\gamma_3$  the inverse image of this path by the homotopy equivalence  $h_{p(A_i)}$ .

We denote by  $[A_1, A_i]$  the path  $\gamma_1 * \gamma_2 * \gamma_3$ .

Now we can define a map  $\sigma : C_i \cap D_j \rightarrow E^n$  by putting  $\sigma(A_1, \dots, A_n) = ([A_1, A_1], \dots, [A_1, A_n])$  for any  $(A_1, \dots, A_n) \in C_i \cap D_j$ .

This map  $\sigma$  is obviously a continuous section of  $e_n^E$ . Applying the Lemma 3.2 we have  $TC_n(E) \leq p + q - 1 = TC_n(X) + TC_n(Y) - 1$ .

**Remark 3.1.** *Suppose that  $X, Y$  are path-connected CW-complexes. Put  $E = X \times Y$  and the map  $p : E \rightarrow X$  to be the projection in the first component. The map  $s : X \rightarrow E$  defined by  $s(x) = (x, y_0)$  is obviously a section of  $p$  and  $h_x : p^{-1}(x) \rightarrow Y$ ,  $h_x(x, y) = y$  is a homotopy equivalence. Then, all the assumptions of Theorem 3.1 are satisfied. And we get again the product inequality (2.1) as stated in Proposition 2.4.*

Theorem 3.1 has been used to compute the higher topological complexity of configuration spaces on some topological manifolds in [3]. We briefly recall the case of configuration space  $F(\mathbb{T}, k)$  on the two-dimensional torus.

The configuration space of  $k$  distinct ordered points in  $\mathbb{T}$  is a subset of  $\mathbb{T}^k$ , defined by

$$F(\mathbb{T}, k) = \{(x_1, \dots, x_k) \in \mathbb{T}^k \mid x_i \neq x_j \text{ with } 1 \leq i \neq j \leq k\}.$$

**Proposition 3.2.** *Let  $k$  be an integer with  $k \geq 2$ . The higher topological complexity of the configuration space of  $k$  distinct ordered points on the 2-dimensional torus  $\mathbb{T}$  is*

$$TC_n(F(\mathbb{T}, k)) = n(k + 1) - 1.$$

Observe that the projection on the first  $k - 1$  coordinates  $\pi_k : F(\mathbb{T}, k) \rightarrow F(\mathbb{T}, k - 1)$  is a fibration with the fiber homotopic to the bouquet of  $k$  circles  $Y_k = \underbrace{\mathbb{S}^1 \vee \dots \vee \mathbb{S}^1}_k$ . It is proved in [3] that this fibration  $\pi_k$  admits a section  $\sigma_k$  for any  $k = 1, \dots$  and satisfies all assumptions of Theorem 3.1. Combining properties of the higher topological complexity mentioned in the previous section and Theorem 3.1 we get

$$TC_n(F(\mathbb{T}, k)) \leq TC_n(F(\mathbb{T}, k - 1)) + TC_n(Y_k) - 1.$$

Moreover, it follows from [3, Theorem 3] that  $TC_n(Y_k) \leq n + 1$  for  $k \geq 2$ . So,

$$TC_n(F(\mathbb{T}, k)) \leq TC_n(F(\mathbb{T}, k - 1)) + n \text{ for all } k \geq 2.$$

By induction on  $k$ , we get

$$TC_n(F_n(\mathbb{T}, k)) \leq TC_n(F(\mathbb{T}, 1)) + n(k - 1) = 2n - 1 + n(k - 1) = n(k + 1) - 1.$$

To prove that  $n(k + 1) - 1$  is the lower bound of  $TC_n(F(\mathbb{T}, k))$  we need to use the lower bound stated in Proposition 2.3 and spectral sequence arguments. We will skip it here. A detailed proof can be found in [3]

## References

- [1] **I. Basabe, J. González, Y.B. Rudyak, and D. Tamaki**, *Higher topological complexity and homotopy dimension of configuration spaces on spheres*, *Algebr. Geom. Topol.* 14 (2014), 2103-2124.
- [2] **Nguyen Viet Dung and Nguyen Van Ninh**, *The Higher Topological Complexity of Complement of Fiber Type Arrangements*, *Acta Math Vietnam* 42, 2(2017), p.249-256.
- [3] **Nguyen Viet Dung and Nguyen Van Ninh**, *The higher topological complexity of configuration spaces of odd-dimensional spheres*, *Singularities - Kagoshima, 2017*, World Scientific (2020). pp 167-183.



- [4] **M.Farber**, *Topological complexity of motion planning*, Discrete Comput. Geom 29(2003) 211 - 221.
- [5] **J.González and M. Grant**, *Sequential motion planning of non-colliding particles in Euclidean spaces*, Proc. Amer. Math. Soc. 143 (2015), 4503-4512.
- [6] **J.González and B. Gutiérrez**, *Topological complexity of collision-free multi-tasking motion planning on orientable surfaces*, Contemp. Math., Amer. Math. Soc. 702 (2018), 151-163.
- [7] **Y.B.Rudiyak**, *On higher analogs of topological complexity*, Topology and its Application, 157(2010) 916-920.
- [8] **A.S.Schwarz**, *The genus of fiber space*, Amer. Math. Sci, Transl 55(1966) 49 - 140.

**Nguyen Viet Dung**

Institute of Mathematics, VAST

Hanoi

Vietnam

vietdung@math.ac.vn; nvdung@cmc-u.edu.vn

**Nguyen Van Ninh**

Thai Nguyen University of Education

Thai Nguyen

Vietnam

nguyenninhsp@gmail.com